


# Characterization of nucleolar localization and exclusion signals in terminal deoxynucleotidyltransferase interacting factor 2/estrogen receptor -binding protein

著者	Fukada Takasi, Shibata Shun, Ueda Toshihiro, Sasaki Katsuhiko, Shimoida Yukiko, Senda-Murata Kaori, Sugimoto Kenji
journal or publication title	Bioscience, Biotechnology, and Biochemistry
volume	83
number	7
page range	1255-1262
year	2019-03-24
URL	<a href="http://hdl.handle.net/10466/00016833">http://hdl.handle.net/10466/00016833</a>

doi: 10.1080/09168451.2019.1591265(<https://doi.org/10.1080/09168451.2019.1591265>)



(b)

	Number of cells with nucleolar localization (%)	Fluorescence Intensity (Nucleolus/ Nucleoplasm)	
mCherry	0%	1.15	
mCherry-NLS2	100%	1.48	
mCherry-NLS3	100%	2.17	
mCherry-NLS2-NLS3	100%	2.34	
mCherry-NLS4	100%	1.61	
mCherry-NLS6	100%	2.13	
mCherry-NLS4-NLS6	100%	2.70	
mCherry-NoES	0%	0.61	
mCherry-101-NLS2-NLS3-134	87%	1.21	
mCherry-101-NLS2-NLS3-134-NoES	0%	0.79	
mCherry-NLS4-NoES	100%	1.34	
mCherry-NoES-NLS6	100%	1.29	
mCherry-NLS4-NoES-NLS6	100%	1.69	