

Roles of Na⁺/Ca²⁺ exchanger isoforms NCX1 and NCX2 in motility in mouse ileum

著者	Nishiyama Kazuhiro, Azuma Yasu-Taka, Morioka Ai, Yoshida Natsuho, Teramoto Midori, Tanioka Kohta, Kita Satomi, Hayashi Satomi, Nakajima Hidemitsu, Iwamoto Takahiro, Takeuchi Tadayoshi
journal or publication title	Naunyn-Schmiedeberg's Archives of Pharmacology
volume	389
number	10
page range	1081-1090
year	2016-10
権利	This is a post-peer-review, pre-copyedit version of an article published in Naunyn-Schmiedeberg's Archives of Pharmacology. The final authenticated version is available online at: http://dx.doi.org/10.1007/s00210-016-1271-1 .
URL	http://hdl.handle.net/10466/15733

doi: 10.1007/s00210-016-1271-1

1
2 **Roles of Na⁺/Ca²⁺ exchanger isoforms NCX1 and NCX2 in motility in mouse ileum**
3
4
5
6
7

8 Kazuhiro Nishiyama • Yasu-Taka Azuma* • Ai Morioka • Natsuho Yoshida • Midori Teramoto •
9

10
11 Kohta Tanioka • Satomi Kita • Satomi Hayashi • Hidemitsu Nakajima • Takahiro Iwamoto •
12

13
14 Tadayoshi Takeuchi
15
16
17
18
19

20
21 K. Nishiyama • Y.T. Azuma • A. Morioka • N. Yoshida • M. Teramoto • K. Tanioka • S. Hayashi •
22

23
24 H. Nakajima • T. Takeuchi
25
26

27 Laboratory of Veterinary Pharmacology, Division of Veterinary Science, Osaka Prefecture
28

29
30 University Graduate School of Life and Environmental Science, Osaka 598-8531, Japan
31
32
33
34
35

36 S. Kita • T. Iwamoto
37
38

39 Department of Pharmacology, Faculty of Medicine, Fukuoka University, Fukuoka 814-0180,
40

41
42 Japan
43
44
45
46
47

48
49 *Corresponding author at: Laboratory of Veterinary Pharmacology, Division of Veterinary
50

51
52 Science, Osaka Prefecture University Graduate School of Life and Environmental Science,
53

54
55 1-58 Rinku-ohraikita, Izumisano, Osaka 598-8531, Japan. TEL/Fax +81-72-463-5264.
56

57
58 E-mail address: azuma@vet.osakafu-u.ac.jp (Y.T. Azuma).
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

The Na⁺/Ca²⁺ exchanger (NCX) is a plasma membrane transporter that is involved in regulating intracellular Ca²⁺ concentrations in various tissues. The physiological roles by which NCX influences gastrointestinal motility are incompletely understood, although its role in the heart, brain and kidney has been widely investigated. In this study, we focused on the functions of the NCX isoforms, NCX1 and NCX2, in the motility of the ileum in the gastrointestinal tract. We investigated the response to electric field stimulation (EFS) in the longitudinal smooth muscle of the ileum obtained from wild-type mice (WT), NCX1-heterozygote knockout mice (NCX1 HET), NCX2 HET and smooth muscle-specific NCX1.3 transgenic mice (NCX1.3 Tg). EFS induced a phasic contraction that persisted during EFS and a tonic contraction that occurred after the end of EFS. We found that the amplitudes of the phasic and tonic contractions were significantly smaller in NCX2 HET, but not in NCX1 HET, compared to WT. Moreover, the magnitudes of acetylcholine (ACh)- and substance P (SP)-induced contractions of NCX2 HET, but not of NCX1 HET, were smaller compared to WT. In contrast, the amplitudes of the phasic and tonic contractions were greater in NCX1.3 Tg compared to WT. Similar to EFS, the magnitude of ACh-induced contraction was greater in NCX1.3 Tg than in WT. Taken together, our findings indicated that NCX1 and NCX2 play important roles in ileal motility and suggest that NCX1 and NCX2 regulate the

1
2 motility in the ileum by controlling the sensitivity of smooth muscles to ACh and SP.
3
4
5
6
7

8 **Key words**
9

10
11 $\text{Na}^+/\text{Ca}^{2+}$ exchanger, ileum, longitudinal smooth muscles, contraction, acetylcholine
12
13
14
15
16

17 **Abbreviations**
18

19
20
21 ACh Acetylcholine
22

23
24 EFS electrical field stimulation
25

26
27 HET heterozygous mice
28

29
30 ICC interstitial cells of Cajal
31

32
33 NCX $\text{Na}^+/\text{Ca}^{2+}$ exchanger
34
35

36
37 SP substance P
38

39
40 NCX1.3 Tg smooth muscle-specific NCX1.3 transgenic mice
41

42
43 WT wild-type
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Introduction

Contractions of longitudinal smooth muscles in the mouse ileum are classified as phasic or tonic. Upon electrical field stimulation (EFS), a phasic contraction exhibits a transient and rapid contraction, whereas a tonic contraction consists of a sustained contraction. The different patterns of phasic and tonic contractions can be explained by the differences in their excitatory transmitters, such as acetylcholine (ACh) and substance P (SP), respectively, followed by the activation of their G protein-coupled receptors. Activated receptors result in the contraction of smooth muscles via Ca^{2+} influx. In gastrointestinal smooth muscles, Ca^{2+} influx triggers Ca^{2+} release from the sarcoplasmic reticulum. The combination of Ca^{2+} influx from extra medium and release from sarcoplasmic reticulum increases the free intracellular Ca^{2+} concentration, thereby allowing Ca^{2+} to bind to a calcium binding protein, which then activates myosin light chain kinase (Berridge, 2008; Wang et al., 2008). Over time, intracellular Ca^{2+} is transported out of the cytosol via pathways involving plasma membrane Ca^{2+} -ATPase, the $\text{Na}^+/\text{Ca}^{2+}$ exchanger (NCX), sarco/endoplasmic reticulum Ca^{2+} -ATPase and mitochondrial Ca^{2+} transporters (Wray and Burdyga, 2010; Webb, 2003). However, the potential role of NCX in gastrointestinal tissue is still incompletely understood among these Ca^{2+} transport pathways.

1
2 Our group focused on the effect of Ca^{2+} movement via NCX on gastrointestinal
3
4 motility because Ca^{2+} homeostasis is central to the regulation of gastrointestinal smooth
5
6 muscle functions. NCX is a plasma membrane transporter that is involved in the regulation of
7
8 intracellular Ca^{2+} concentrations in tissues, such as the brain, kidney, and smooth muscle.
9
10
11
12
13
14 NCX electrogenically exchanges Na^+ and Ca^{2+} across the plasma membrane, depending on the
15
16
17 membrane potential and transmembrane gradients. The mammalian NCX family consists of
18
19
20 three isoforms: NCX1 (Nicoll et al., 1990), NCX2 (Li et al., 1994) and NCX3 (Nicoll et al.,
21
22
23 1996). To specifically assess the physiological role of NCX isoforms, we used NCX1
24
25
26 heterozygote knockout mice (HET) (Wakimoto et al., 2000) and NCX2 HET (Gotoh et al.,
27
28
29 2015). We previously demonstrated that NCX2 HET, but not NCX1 HET, displays altered
30
31
32 motility in the distal colon (Azuma et al., 2012). In addition, we recently demonstrated that
33
34
35 NCX1 HET and NCX2 HET display altered motility in the gastric fundus (Azuma et al.,
36
37
38 2016a). Currently, little is known about the potential role of NCX isoforms in small intestinal
39
40
41 motility. This study aimed to determine the effect of NCX isoforms heterozygous deficiency
42
43
44 on the motility of the ileum. To further test whether smooth muscle NCX1 plays a critical role
45
46
47 in motility in the ileum, we used transgenic mice expressing canine NCX1.3 driven by the
48
49
50 smooth muscle-specific α -actin promoter. To better understand the action of NCX isoforms,
51
52
53 we used an organ-tissue bath system to characterize the motility of longitudinal muscle strips
54
55
56
57
58 isolated from the ileum.
59
60
61
62
63
64
65

Materials and Methods

Drugs

ACh and atropine were purchased from Wako Pure Chemical (Osaka, Japan). An Alexa Fluor 488-labeled goat anti-rabbit IgG was purchased from Molecular Probes Inc. (Eugene, OR, USA). A mouse polyclonal antibody against PGP9.5 was purchased from UltraClone Limited (Isle of Wight, UK). SP was purchased from the Peptide Institute (Osaka, Japan). Rabbit polyclonal antibodies against NCX1 and NCX2 were produced as previously described (Iwamoto et al., 1998).

Animals

NCX1 HET, NCX2 HET and smooth muscle-specific NCX1.3 transgenic mice (NCX1.3 Tg) were produced as reported previously (Wakimoto et al., 2000; Gotoh et al., 2015; Iwamoto et al., 2004). These three mice on the C57BL/6 background appeared healthy and were comparable in all analyses to age-matched wild-type mice (WT). All procedures used in this study were performed according to the institutional policies of the Osaka Prefecture University Animal Care and Use Committee.

Immunofluorescent staining

1
2 Immunofluorescent staining on frozen sections was performed as previously described
3
4
5 (Azuma et al., 2008) with some modifications (Nishiyama et al., 2013). Immunoreactivity of
6
7
8 NCX1 and NCX2 was detected using an Alexa Fluor 488-labeled goat anti-rabbit IgG
9
10
11 antibody. To detect neurons, a mouse antibody against PGP9.5 was used, and the
12
13
14 immunoreactivity of PGP9.5 was detected using an Alexa Fluor 488-labeled goat anti-mouse
15
16
17 IgG antibody. Confocal images were obtained using a laser-scanning microscope (C1si; Nikon
18
19
20 Corporation, Tokyo).
21

22 23 24 25 26 27 Quantitative real-time PCR

28
29
30 Quantitative real-time PCR for NCX1 and NCX2 mRNA was performed as described
31
32
33 previously (Azuma et al., 2010a) with some modifications (Matsuo et al., 2015). Briefly, the
34
35
36 ileum was removed from the mice. Total RNA was extracted. The primers used for the
37
38
39 amplification of NCX1 and NCX2 were as follows: 5'-CGCTGGGGAAGATGACGATG -3'
40
41
42 and 5'-TGGGACGAAGGCAAACAGAAC -3', and 5' -ATGGCTCCCTTGGCTTTGATG -3 '
43
44
45 and 5' -CAGCGGTAGGAACCTTGGC -3 ', respectively. The amplification of HPRT mRNA
46
47
48 was used as an endogenous control for each experimental sample to account for differences in
49
50
51 the amount and quality of total RNA added to each reaction.
52

53 54 55 56 57 58 Western blot analysis

59
60
61
62
63
64
65

1 Western blot analysis was performed using a previously described method (Azuma et al.,
2 [2010b](#)) with some modifications (Nishiyama et al., [2014](#)). Briefly, the ileum was harvested
3
4
5 with Cell Lysis Buffer (Cell Signaling). Aliquots were electrophoresed on a 10%
6
7
8 polyacrylamide gel and subsequently blotted on a PVDF membrane. The PVDF membrane
9
10
11 was reacted with antibodies against NCX1 or NCX2 and then reacted with a
12
13
14 peroxidase-conjugated secondary antibody. Finally, the proteins that reacted with the antibody
15
16
17 were visualized using ECLTM detection reagents (GE Healthcare).
18
19
20
21
22
23
24
25
26

27 Recording of responses to EFS in longitudinal smooth muscles of the ileum

28
29 Responses to EFS were recorded using previously described methods (Tekeuchi et al., [2007](#);
30
31
32 Azuma et al., [2016](#)). Briefly, the ileum was removed from mice (10–15 weeks old).
33
34
35 Whole-wall strips were prepared in the orientation of the longitudinal muscle layer. These
36
37
38 strips were mounted in organ baths and held at a resting tension of 0.5 g. Responses to EFS
39
40
41 were detected using isotonic force transducers (TD-112A; Nihonkohden, Tokyo, Japan).
42
43
44 Specifically, the strips were exposed to EFS with trains of 100 pulses of 0.5 ms and 30 V for
45
46
47 10 sec, with a 10-min interval between tests. Atropine (1 μ M) was directly added to the bath at
48
49
50
51 least 10 min prior to EFS. Phasic and tonic contractions were analyzed by measuring the
52
53
54 extent of the maximal contraction in response to 60 mM KCl.
55
56
57
58
59
60
61
62
63
64
65

Statistical analysis

The results were expressed as the mean \pm S.E. Statistical significance was determined using one-way ANOVA for non-repeated measures to detect differences among WT, NCX1 HET and NCX2 HET. The differences between groups were determined using the Tukey-Kramer test. The statistical significance of the parametric data was evaluated using a two-tailed Student's t-test to detect differences between WT and NCX1.3 Tg. A p value less than 0.05 was considered significant.

Results

Localization of NCX1 and NCX2

We initially investigated the expression and localization of NCX1 and NCX2 in the ileum of WT using immunofluorescent staining. As shown in Figure 1A, strong immunoreactivity of NCX1 and NCX2 was observed within the myenteric plexus layers. The expression of NCX1 and NCX2 was also observed in the longitudinal and circular muscle layers. To estimate the histological change of NCX1- and NCX2-HET tissue preparations in the ileum, we dyed myenteric neurons with the PGP9.5 antibody. As shown in Figure 1B, immunoreactivity of PGP9.5 was observed within the myenteric plexus layers among WT, NCX1 HET and NCX2 HET. In addition, there are no marked histological changes in the ileum between WT and NCXs HET.

Decreased expression of NCX1 and NCX2

As shown in Figure 2A, the mRNA expression level of NCX1 was significantly lower in NCX1 HET, whereas NCX2 HET expressed significantly lower levels of NCX2 mRNA. In accordance with these results, the protein expression level of NCX1 was significantly lower in NCX1 HET, whereas NCX2 HET expressed significantly lower levels of NCX2 protein (Fig. 2B).

EFS-induced responses

First, we evaluated the amplitude and spike numbers of spontaneous contractions. Consequently, there were no significant changes among WT, NCX1 HET and NCX2 HET (data not shown). Next, we investigated the frequency-response relationship of EFS at 1, 3 and 10 Hz in the ileum from WT, NCX1 HET and NCX2 HET. Figure 3A shows representative recording traces of responses to EFS. In WT, EFS induced both a phasic contraction that persisted during the stimulus and a tonic contraction that was recorded after the end of the stimulus. To measure the extent of the phasic and tonic contractions, we analyzed the extent of the maximal contractions in response to 60 mM KCl among WT, NCX1 HET and NCX2 HET. The magnitudes of maximal contractions in response to 60 mM KCl were similar among WT, NCX1 HET and NCX2 HET (Fig. 4). Importantly, the magnitude of the phasic

1 contraction at 1 Hz was smaller in NCX2 HET, but not NCX1 HET, compared to WT (Fig. 3B
2
3
4 top). However, NCX1 HET and NCX2 HET showed phasic contractions at 3 Hz and 10 Hz in
5
6
7 a manner similar to WT (Fig. 3B top). In contrast, the tonic contractions at 1 and 3 Hz were
8
9 smaller only in NCX2 HET (Fig. 3B bottom).
10
11
12

13
14 We previously demonstrated that ACh is the representative transmitter of phasic
15
16 contractions and that SP is the main transmitter of tonic contractions in the mouse ileum
17
18 (Takeuchi et al., 2007). To characterize the tonic contraction, EFS was performed after the
19
20 tissues were incubated with atropine, a muscarinic ACh receptor antagonist. The magnitude of
21
22 the phasic contraction was completely suppressed in WT, NCX1 HET and NCX2 HET.
23
24 Nevertheless, tonic contractions remained. The magnitude of the tonic contraction was smaller
25
26 in NCX2 HET, but not in NCX1 HET, compared to WT (Fig. 3C). Taken together, these
27
28 results indicate that NCX2 HET produced decreases in the response to excitatory mediators,
29
30 such as ACh and SP.
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 Responses of smooth muscles

46
47 Because NCX1 and NCX2 are expressed in longitudinal smooth muscles and in neurons of the
48
49 myenteric plexus layers (Fig. 1A), we determined the effect of NCX isoforms in heterozygous
50
51 deficient mice on the contractile response to ACh and SP in smooth muscle cells. We
52
53 examined the responses to three different concentrations of ACh and two different
54
55
56
57
58
59
60
61
62
63
64
65

1 concentrations of SP. ACh and SP induced contractions in a dose-dependent manner (Fig. 5
2
3
4
5 and Fig. 6). The magnitude of the ACh-induced contractions was smaller in NCX2 HET than
6
7
8 in WT when ACh concentration was only 0.1 μ M (Fig. 5B). Similarly, the magnitudes of the
9
10
11 SP-induced contractions were smaller in NCX2 HET than in WT at both concentrations
12
13
14 examined (Fig. 6B). However, ACh and SP induced similar magnitudes of contraction in
15
16
17 NCX1 HET compared to WT (Fig. 5B and Fig. 6B). Thus, NCX2 heterozygous deficiency in
18
19
20 smooth muscles may lower the sensitivity of these cells to ACh and SP.
21
22
23
24
25
26

27 Overexpression of NCX1

28
29 These observations suggested that NCX1 is not likely to affect ileum contractions, although
30
31
32 NCX1 was expressed in longitudinal and circular muscle layers and in the myenteric plexus
33
34
35 layers. To further test whether NCX1 plays a critical role in ileal contractions, we examined
36
37
38 the EFS-induced responses in the longitudinal smooth muscles of the ileum in NCX1.3 Tg in
39
40
41 the next series of experiments. We confirmed the overexpression of NCX1 in the ileum of
42
43
44 NCX1.3 Tg using western blot analysis. As shown in Figure 7, NCX1 protein was
45
46
47 overexpressed in the ileum of NCX1.3 Tg at a two-fold higher level compared to endogenous
48
49
50
51 NCX1.
52
53
54
55
56
57

58 EFS-induced responses in NCX1.3 Tg

59
60
61
62
63
64
65

1
2 To measure the extent of the phasic and tonic contractions, we analyzed the extent of the
3
4 maximal contractions in response to 60 mM KCl between WT and NCX1.3 Tg. The
5
6 magnitudes of the maximal contractions in response to 60 mM KCl were similar between WT
7
8 and NCX1.3 Tg (Fig. 8). We found that the magnitudes of the phasic and tonic contractions
9
10 were greater in NCX1.3 Tg than in WT at 10Hz (Fig. 9B). Tonic contractions during atropine
11
12 administration were also greater in NCX1.3 Tg than in WT at 10Hz (Fig. 9D). The time from
13
14 full amplitudes to half amplitudes of the phasic contraction were compared in WT and
15
16 NCX1.3 Tg (Fig. 9A). Interestingly, the time to half amplitudes was approximately 4 seconds
17
18 longer in NCX1.3 Tg than in WT (Fig. 9C).
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33 Responses of smooth muscle cells in NCX1.3 Tg

34
35 Consistent with the results that phasic contraction was greater in NCX1.3 Tg than in WT, the
36
37 magnitude of ACh-induced contractions was greater in NCX1.3 Tg than in WT (Fig. 10A). In
38
39 contrast, NCX1.3 Tg exhibited SP-induced contractions that were similar to those of WT (Fig.
40
41 10C). The time from full amplitudes to half amplitudes of the ACh and SP-induced
42
43 contractions were significantly longer in NCX1.3 Tg than in WT (Fig. 10B and 10D).
44
45
46
47
48
49
50
51
52
53
54

55 Discussion

56
57
58 The first aim of this study was to investigate the physiological role of NCX isoforms in the
59
60
61
62
63
64
65

1 motility of the ileum. Our immunohistochemical results provide evidence indicating the
2
3
4
5 presence of NCX1- and NCX2-positive cells in the myenteric plexus in addition to the
6
7
8 longitudinal and circular muscle layers. Our findings indicate a role for the NCX isoform in
9
10
11 regulating murine ileal motility and highlight the importance of NCX isoforms in this
12
13
14 regulation. We demonstrated that NCX2 HET show decreased amplitudes of EFS-induced
15
16
17 contractions. This decreased contraction is not pronounced in NCX1 HET, which suggests that
18
19
20 NCX2 plays a dominant role than NCX1 in ileal motility. How does NCX2 HET decrease the
21
22
23 contractions? There are at least two potential explanations. One possibility may be a
24
25
26 decreased release of an excitatory transmitter, such as ACh or SP, from myenteric neurons
27
28
29 during EFS. The second possibility is an attenuated sensitivity to the excitatory transmitters of
30
31
32 smooth muscles. In the present study, the amplitude of the ACh-induced contraction was
33
34
35 significantly decreased in NCX2 HET compared to WT, suggesting that decreased phasic
36
37
38 contraction in NCX2 HET was associated with the attenuated sensitivity of smooth muscles to
39
40
41 ACh. Similar to ACh, the amplitude of SP-induced contraction was significantly decreased in
42
43
44 NCX2 HET compared to WT, suggesting that the decreased tonic contraction in NCX2 HET is
45
46
47 associated with the attenuated sensitivity of smooth muscles to SP. Consistent with previous
48
49
50 findings, we found that there are NCX2-positive cells in the myenteric plexus. What is the
51
52
53 role of NCX2 expressed in the myenteric plexus? It is important to demonstrate which cell
54
55
56
57
58 type contributes to the regulation of motility in the ileum. We hypothesized that the decreased
59
60
61
62
63
64
65

1 expression of NCX2 in the smooth muscles of NCX2 HET would result in small contractions.
2
3
4
5 However, to date, the role of NCX2 in the myenteric plexus of the ileum remains unclear. We
6
7
8 previously showed that ACh release decreased in the colon of NCX2 HET (Azuma et al.,
9
10 [2012](#)). Thus, the possibility that a decrease in the release of the transmitters may also be
11
12 responsible for the small EFS-induced contractions in the ileum of NCX2 HET cannot be ruled
13
14 out. Further experiments are required to investigate the direct role of NCX2 in the myenteric
15
16 plexus. Further investigation will focus on precisely distinguishing between the contributions
17
18 of myenteric plexus NCX2 versus smooth muscle NCX2 to the contractile function of the
19
20 ileum. In smooth muscles, elevated intracellular Ca^{2+} concentrations are essential for the
21
22 contraction in response to ACh or SP. In all of the previous and recent findings, NCX,
23
24 including NCX2, act to increase the efflux of Ca^{2+} and influx of Na^+ , resulting in a decrease in
25
26 smooth muscle tone. Originally, we expected that the decrease in the expression level of
27
28 NCX2 in NCX2 HET might result in an increase in smooth muscle tone, which is inconsistent
29
30 with the present results. If NCX2 contributes to Ca^{2+} efflux, then NCX2 HET may result in
31
32 large contractions. However, the present results revealed that NCX2 HET showed small
33
34 contractions. Thus, we hypothesized that NCX2 contributes to Ca^{2+} influx, but not Ca^{2+} efflux,
35
36 during contraction in response to ACh or SP. It is likely that NCX2 HET occurs following
37
38 weakly elevated intracellular Ca^{2+} concentrations. In support of this hypothesis, it has been
39
40 reported that Ca^{2+} influx via NCX contributes to neurotransmitter release in rat cortical
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 neurons (Wu et al., 2008). Furthermore, the promotion of Ca^{2+} influx via NCX has been
2
3
4 suggested to occur during heart failure (Sato et al., 2000) and ischemia (Imahashi et al.,
5
6
7 2005). Together with previous evidence, these results suggest that NCX2 may contribute to
8
9
10 Ca^{2+} influx in the ileum.
11

12
13
14 Furthermore, this study indicated that 60 mM KCl-induced contractions were similar
15
16
17 among WT, NCX1 HET and NCX2 HET, and between WT and NCX1.3 Tg. If NCX1 and
18
19
20 NCX2 can contribute to Ca^{2+} influx in the ileum, why were 60 mM KCl-induced contractions
21
22
23 in NCXs HET and NCX1.3 Tg not changed? It is likely that the role and contribution of Ca^{2+}
24
25
26 influx is different between ACh- and SP-induced contractions and KCl-induced contraction.
27
28
29 Another possibility is that the role and contribution of NCX is different between ACh- and
30
31
32 SP-induced contractions and KCl-induced contraction. A further issue that is raised by the
33
34
35 current work is to determine the role of NCX in the contraction machinery of the ileum.
36
37
38

39
40 Regarding NCX1 HET, our observations suggested that NCX1 is not likely to affect
41
42
43 ileal contraction, although NCX1 was expressed in both the longitudinal and circular muscle
44
45
46 layers and in the myenteric plexus layers. Then, what is the role of NCX1? Using NCX1.3 Tg,
47
48
49 we demonstrated that NCX1 overexpression enhanced EFS-induced contractions. Importantly,
50
51
52 findings using NCX1.3 Tg indicated a prolonged time period from full amplitudes to half
53
54
55 amplitudes. We found that NCX1 overexpression also enhanced ACh-induced contraction,
56
57
58 which suggests that the enhanced contraction in NCX1.3 Tg is associated with an increased
59
60
61
62
63
64
65

1
2 sensitivity to ACh. Next, we considered the mechanism by which NCX1.3 Tg showed
3
4
5 enhanced contraction and prolonged time to half amplitudes. Elevated intracellular Ca^{2+}
6
7
8 concentrations are essential for contractions in smooth muscles. In urinary bladder smooth
9
10
11 muscles, NCX1.3 Tg showed enhanced contractions and a prolonged duration of Ca^{2+} sparks
12
13
14 (Yamamura et al., 2013). In combination with previous findings, NCX1.3 Tg has been
15
16
17 hypothesized to experience increased Ca^{2+} influx via NCX, which extends the period of high
18
19
20 Ca^{2+} levels and maintains the contraction. Consistent with this hypothesis, NCX1
21
22
23 overexpression also prolonged the time to half amplitudes, though it had no effect on the
24
25
26 amplitude of the SP-induced contractions. Taken together, these results suggest that increased
27
28
29 Ca^{2+} influx via NCX after the contraction extends the period of high Ca^{2+} levels, followed by a
30
31
32 prolonged time to half amplitudes.
33

34
35
36 Given that NCX1 and NCX2 proteins are normally expressed on all cell types in the
37
38
39 gastrointestinal tissues-nerves, glia, interstitial cells of Cajal (ICC) and smooth muscle (Oda
40
41
42 et al., 2011; Ota et al., 2013; Yamakawa T et al., 2012), we predicted that disruption of
43
44
45 $\text{Na}^+/\text{Ca}^{2+}$ homeostasis in NCX1 HET and NCX2 HET may affect many signaling pathways in
46
47
48 all or any of the cells that contribute to gastric motility. ICC, which are also called pacemaker
49
50
51 cells (Kito and Suzuki, 2003; Kito et al., 2005), have been suggested to be involved in the
52
53
54 neurotransmission in gastrointestinal smooth muscles (Bayguinov et al., 2010; Tanahashi et
55
56
57 al., 2014). In addition, ICC-MY have been suggested to express NCX1 and to be involved in
58
59
60
61
62
63
64
65

1
2 its Ca^{2+} handling (Cho and Daniel, 2005; Lowie et al., 2011). Thus, these results can
3
4
5 potentially imply the deficient function of ICC in NCXs HET and the function of ICC in
6
7
8 NCX1.3 Tg. A further issue that was raised by the current work was to determine the role of
9
10
11 NCX expressed in ICC.
12

13
14 In conclusion, our findings suggest that NCX isoforms play prominent roles in
15
16
17 contraction in the ileum. Thus, NCXs HET and NCX1.3 Tg may be valuable tools for
18
19
20 delineating the mechanisms of NCX isoforms action in gastrointestinal motility. To this end,
21
22
23
24 the current study may provide useful information for the identification of therapeutic targets.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- Azuma YT, Hayashi S, Nishiyama K, Kita S, Mukai K, Nakajima H, Iwamoto T, Takeuchi T (2016a) $\text{Na}^+/\text{Ca}^{2+}$ exchanger-heterozygote knockout mice display increased relaxation in gastric fundus and accelerated gastric transit in vivo. *Neurogastroenterol Motil* in press.
- Azuma YT, Hagi K, Shintani N, Kuwamura M, Nakajima H, Hashimoto H, Baba A, Takeuchi T (2008) PACAP provides colonic protection against dextran sodium sulfate induced colitis. *J Cell Physiol* 216:111-119
- Azuma YT, Samezawa N, Nishiyama K, Nakajima H, Takeuchi T (2016b) Differences in time to peak carbachol-induced contractions between circular and longitudinal smooth muscles of mouse ileum. *Naunyn Schmiedebergs Arch Pharmacol* 389:63-72
- Azuma YT, Nishiyama K, Kita S, Komuro I, Nakajima H, Iwamoto T, Takeuchi T (2012) $\text{Na}^+/\text{Ca}^{2+}$ exchanger 2-heterozygote knockout mice display decreased acetylcholine release and altered colonic motility in vivo. *Neurogastroenterol Motil* 24:e600-e610
- Azuma YT, Matsuo Y, Kuwamura M, Yancopoulos GD, Valenzuela DM, Murphy AJ, Nakajima H, Karow M, Takeuchi T (2010b) Interleukin-19 protects mice from innate-mediated colonic inflammation. *Inflamm Bowel Dis* 16:1017–1028

- 1
2 Azuma YT, Nishiyama K, Matsuo Y, Kuwamura M, Morioka A, Nakajima H, Takeuchi T
3
4
5 (2010a) PPAR α contributes to colonic protection in mice with DSS-induced colitis. *Int*
6
7
8 *Immunopharmacol* 10:1261-1267
9
- 10
11 Bayguinov PO, Hennig GW, Smith TK (2010) Ca²⁺ imaging of activity in ICC-MY during
12
13
14 local mucosal reflexes and the colonic migrating motor complex in the murine large
15
16
17 intestine. *J Physiol* 588:4453-4474.
18
19
- 20
21 Berridge MJ (2008) Smooth muscle cell calcium activation mechanisms. *J Physiol*
22
23
24 586:5047-5061
25
26
- 27 Cho WJ, Daniel EE (2005) Proteins of interstitial cells of Cajal and intestinal smooth muscle,
28
29
30 colocalized with caveolin-1. *Am J Physiol Gastrointest Liver Physiol* 288:G571-G585.
31
32
- 33 Gotoh Y, Kita S, Fujii M, Tagashira H, Horie I, Arai Y, Uchida S, Iwamoto T (2015) Genetic
34
35
36 knockout and pharmacologic inhibition of NCX2 cause natriuresis and hypercalciuria.
37
38
39 *Biochem Biophys Res Commun* 456:670-675
40
41
- 42 Imahashi K, Pott C, Goldhaber JI, Steenbergen C, Philipson KD, Murphy E (2005)
43
44
45 Cardiac-specific ablation of the Na⁺-Ca²⁺ exchanger confers protection against
46
47
48 ischemia/reperfusion injury. *Circ Res* 97:916-921.
49
50
- 51
52 Iwamoto T, Kita S, Zhang J, Blaustein MP, Arai Y, Yoshida S, Wakimoto K, Komuro I,
53
54
55 Katsuragi T (2004) Salt-sensitive hypertension is triggered by Ca²⁺ entry via Na⁺/Ca²⁺
56
57
58 exchanger type-1 in vascular smooth muscle. *Nat Med* 10:1193-1199
59
60
61
62
63
64
65

1 Iwamoto T, Pan Y, Nakamura TY, Wakabayashi S, Shigekawa M (1998) Protein kinase
2 C-dependent regulation of Na⁺/Ca²⁺ exchanger isoforms NCX1 and NCX3 does not
3
4
5
6
7
8
9 require their direct phosphorylation. *Biochemistry* 37:17230-17238

10
11 Kito Y, Suzuki H (2003) Electrophysiological properties of gastric pacemaker potentials. *J*
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Kito Y, Ward SM, Sanders KM (2003) Pacemaker potentials generated by interstitial cells of
Cajal in the murine intestine. *Am J Physiol Cell Physiol* 288:C710-C720.

Kito Y, Ward SM, Sanders KM (2005) Pacemaker potentials generated by interstitial cells of
Cajal in the murine intestine. *Am J Physiol Cell Physiol* 288:C710-C720.

Li Z, Matsuoka S, Hryshko LV, Nicoll DA, Bersohn MM, Burke EP, Lifton RP, Philipson KD
(1994) Cloning of the NCX2 isoform of the plasma membrane Na⁺-Ca²⁺ exchanger. *J*
Biol Chem 269:17434-17439

Lowie BJ, Wang XY, White EJ, Huizinga JD (2011) On the origin of rhythmic calcium
transients in the ICC-MP of the mouse small intestine. *Am J Physiol Gastrointest Liver*
Physiol 301:G835-G845.

Matsuo Y, Azuma YT, Kuwamura M, Kuramoto N, Nishiyama K, Yoshida N, Ikeda Y,
Fujimoto Y, Nakajima H, Takeuchi T (2015) Interleukin 19 reduces inflammation in
chemically induced experimental colitis. *Int Immunopharmacol* 29:468-475

Nicoll DA, Longoni S, Philipson KD (1990) Molecular cloning and functional expression of the
cardiac sarcolemmal Na⁺-Ca²⁺ exchanger. *Science* 250:562-565

Nicoll DA, Quednau BD, Qui Z, Xia YR, Lusa AJ, Philipson KD (1996) Cloning of a third

1 mammalian Na⁺-Ca²⁺ exchanger, NCX3. *J Biol Chem* 271:24914-24921

2
3
4
5 Nishiyama K, Morioka A, Kita S, Nakajima H, Iwamoto T, Azuma YT, Takeuchi T (2014)

6
7
8 Na⁺/Ca²⁺ exchanger 1 transgenic mice display increased relaxation in the distal colon.

9
10
11 *Pharmacology* 94:230-238

12
13
14 Nishiyama K, Azuma YT, Kita S, Azuma N, Hayashi S, Nakajima H, Iwamoto T, Takeuchi T

15
16
17 (2013) Na⁺/Ca²⁺ exchanger 1/2 double-heterozygote knockout mice display increased

18
19
20 nitric oxide component and altered colonic motility. *J Pharmacol Sci* 123:235-245

21
22
23 Oda T, Kume T, Izumi Y, Ishihara K, Sugmimoto H, Akaike A (2011) Na⁺/Ca²⁺ exchanger

24
25
26 inhibitors inhibit neurite outgrowth in PC12 cells. *J Pharmacol Sci* 116:128-131.

27
28
29 Ota Y, Kawanai T, Watanabe R, Nishimura A, Ago Y, Takuma K, Matsuda T (2013) Effect of

30
31
32 Overexpression of the Brain-Specific Na⁽⁺⁾/Ca⁽²⁺⁾ Exchanger Splice Variant NCX1.5

33
34
35 on NO Cytotoxicity in HEK293 Cells. *J Pharmacol Sci* 121:351-354.

36
37
38 Satoh H, Ginsburg KS, Qing K, Terada H, Hayashi H, Bers DM (2000) KBR7943 block of

39
40
41 Ca²⁺ influx via Na⁺/Ca²⁺ exchange does not alter twitches or glycoside inotropy but

42
43
44 prevents Ca²⁺ overload in rat ventricular myocytes. *Circulation* 101:1441-1446.

45
46
47 Tanahashi Y, Ichimura Y, Kimura K, Matsuyama H, Iino S, Komori S, Unno T (2014)

48
49
50 Cholinergic neuromuscular transmission mediated by interstitial cells of Cajal in the

51
52
53 myenteric layer in mouse ileal longitudinal smooth muscles. *Naunyn Schmiedebergs*

54
55
56
57
58 *Arch Pharmacol* 387:377-388.

59
60
61
62
63
64
65

- 1
2 Takeuchi T, Tanaka K, Nakajima H, Matsui M, Azuma YT (2007) M2 and M3 muscarinic
3
4
5 receptors are involved in enteric nerve-mediated contraction of the mouse ileum:
6
7
8 Findings obtained with muscarinic-receptor knockout mouse. *Am J Physiol Gastrointest*
9
10
11 *Liver Physiol* 292:G154-G164
12
13
14 Wang Y, Deng X, Hewavitharana T, Soboloff J, Gill DL (2008) Stim, ORAI and TRPC
15
16
17 channels in the control of calcium entry signals in smooth muscle. *Clin Exp Pharmacol*
18
19
20 *Physiol* 35:1127–1133
21
22
23
24 Wakimoto K, Kobayashi K, Kuro-O M, Yao A, Iwamoto T, Yanaka N, Kita S, Nishida A,
25
26
27 Azuma S, Toyoda Y, Omori K, Imahie H, Oka T, Kudoh S, Kohmoto O, Yazaki Y,
28
29
30 Shigekawa M, Imai Y, Nabeshima Y, Komuro I (2000) Targeted disruption of Na⁺/Ca²⁺
31
32
33 exchanger gene leads to cardiomyocyte apoptosis and defects in heartbeat. *J Biol Chem*
34
35
36 275:36991-36998
37
38
39
40 Webb RC (2003) Smooth muscle contraction and relaxation. *Adv Physiol Educ* 27:201-206
41
42
43 Wray S, Burdyga T (2010) Sarcoplasmic reticulum function in smooth muscle. *Physiol Rev*
44
45
46 90:113-178
47
48
49 Wu MP, Kao LS, Liao HT, Pan CY (2008) Reverse mode Na⁺/Ca²⁺ exchangers trigger the
50
51
52 release of Ca²⁺ from intracellular Ca²⁺ stores in cultured rat embryonic cortical neurons.
53
54
55 *Brain Res* 1201:41-51.
56
57
58
59
60
61
62
63
64
65

1
2 Yamakawa T, Watanabe Y, Watanabe H, Kimura J (2012) Inhibitory effect of cibenzoline on
3
4
5 Na(+)/Ca(2+) exchange current in guinea-pig cardiac ventricular myocytes. J Pharmacol
6
7
8 Sci 120:59-62.
9

10
11 Yamamura H, Cole WC, Kita S, Hotta S, Murata H, Suzuki Y, Ohya S, Iwamoto T, Imaizumi
12
13
14 Y (2013) Overactive bladder mediated by accelerated Ca²⁺ influx mode of Na⁺/Ca²⁺
15
16
17 exchanger in smooth muscle. Am J Physiol Cell Physiol 305:C299-C308
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 **Fig. 1.** Immunohistochemical staining. (A) NCX expression in the tissue sections of the ileum
3
4 of WT. Tissue sections were stained for NCX1 or NCX2 (green). Images shown are
5
6 representative of three experiments. CM, circular muscle layer; MP, myenteric plexus layer;
7
8 LM, longitudinal muscle layer. Scale bar, 50 μ m. (B) Immunohistochemical staining of the
9
10 tissue sections in the ileum of NCX HETs. Tissue sections were stained for PGP9.5 (green)
11
12 which is a marker of neurons. Images shown are representative of three experiments. MP,
13
14 myenteric plexus layer. Scale bar, 20 μ m.
15
16
17
18
19
20
21
22
23
24
25
26

27 **Fig. 2.** Decreased expression of NCX in the ileum. (A) The mRNA expression level of NCX1
28
29 in WT (n=7), NCX1 HET (n=7) and NCX2 HET (n=5) and the mRNA expression level of
30
31 NCX2 in WT (n=7), NCX1 HET (n=6) and NCX2 HET (n=7) were examined using
32
33 quantitative real-time PCR. The mRNA level of each NCX is shown as a fold increase relative
34
35 to the level of HPRT mRNA. *P<0.05 compared with WT. (B) Protein expressions of NCX in
36
37 the ileum. Total cell lysates were prepared from the ileum in WT (n=4) and NCX1 HET (n=4)
38
39 and blotted with antibody against NCX1, and in WT (n=7) and NCX2 HET (n=7) and blotted
40
41 with antibody against NCX2. *P<0.05 compared with WT.
42
43
44
45
46
47
48
49
50
51
52
53
54

55 **Fig. 3.** Decreased phasic and tonic contractions in NCX2 HET. EFS-induced contractions in
56
57 longitudinal smooth muscles isolated from the ileum in WT (n=7), NCX1 HET (n=11) and
58
59
60
61
62
63
64
65

1 NCX2 HET (n=7). The muscle strips were stimulated by EFS at 1, 3 and 10 Hz for 10 s. (A)
2
3
4
5 Representative recording traces of EFS-induced contractions are shown. Bars indicate the
6
7
8 duration (10 s) of EFS. After basal tones were recorded, the chart speed was increased to make
9
10
11 the EFS-induced contraction clear. (B) Quantitative data on phasic and tonic contractions.
12
13
14 Phasic and tonic contractions were expressed as percentages of 60 mM KCl-induced
15
16
17 contraction. **P<0.01 for WT vs. NCX2 HET. (C) EFS in the presence of atropine in
18
19
20 longitudinal smooth muscles isolated from the ileum in WT (n=5), NCX1 HET (n=5) and
21
22
23 NCX2 HET (n=4). Quantitative data on the EFS-induced tonic contraction. Tonic contractions
24
25
26 were expressed as percentages of 60 mM KCl-induced contraction. *P<0.05 for WT vs. NCX2
27
28
29 HET.
30

31
32
33
34
35
36 **Fig. 4.** KCl-induced contraction. KCl-induced contractions in longitudinal smooth muscles
37
38
39 isolated from the ileum in WT (n=21), NCX1 HET (n=10) and NCX2 HET (n=9). Quantitative
40
41
42 data on KCl-induced contractions are expressed as mm.
43
44
45
46
47

48
49 **Fig. 5.** ACh-induced contraction. ACh-induced contractions in longitudinal smooth muscles
50
51
52 isolated from the ileum in WT (n=4), NCX1 HET (n=5) and NCX2 HET (n=4). (A)
53
54
55 Representative recording traces of ACh-induced contractions are shown. (B) Quantitative data
56
57
58 on ACh-induced contractions. ACh-induced contractions are expressed as percentages of 60
59
60
61
62
63
64
65

1 mM KCl-induced contractions. **P<0.01 for WT vs. NCX2 HET.

2
3
4
5
6
7
8 **Fig. 6.** SP-induced contraction. SP-induced contractions in longitudinal smooth muscles
9
10 isolated from the ileum in WT (n=4), NCX1 HET (n=5) and NCX2 HET (n=7). (A)
11
12 Representative recording traces of SP-induced contractions are shown. (B) Quantitative data
13
14 on SP-induced contractions. SP-induced contractions are expressed as percentages of 60 mM
15
16 KCl-induced contractions *P<0.05 for WT vs. NCX2 HET.
17
18
19
20
21
22
23
24
25
26

27 **Fig. 7.** NCX1 expression in the ileum. Total cell lysates were prepared from the ileum in WT
28
29 (n=3) and NCX1.3 Tg (n=3) and blotted with antibody against NCX1. *P<0.05 for WT vs.
30
31 NCX1.3 Tg.
32
33
34
35
36
37
38

39 **Fig. 8.** KCl-induced contraction. KCl-induced contractions in longitudinal smooth muscles
40
41 isolated from the ileum in WT (n=13) and NCX1.3 Tg (n=11). Quantitative data on
42
43 KCl-induced contractions are expressed as mm.
44
45
46
47
48
49
50

51 **Fig. 9.** EFS-induced contraction in NCX1.3 Tg. (A-C) EFS-induced contractions in
52
53 longitudinal smooth muscles isolated from the ileum in WT (n=6) and NCX1.3 Tg (n=7). The
54
55 muscle strips were stimulated by EFS at 10 Hz for 10 s. (A) Representative recording traces of
56
57
58
59
60
61
62
63
64
65

1 EFS-induced contraction are shown. Bars indicate the duration (10 s) of EFS. (B) Quantitative
2 data on phasic and tonic contractions. Phasic and tonic contractions were expressed as
3 percentages of 60 mM KCl-induced contraction. *P<0.05, **P<0.01 for WT vs. NCX1.3 Tg.
4
5 (C) t1/2 describes time from full amplitudes to half amplitudes of the phasic contraction.
6
7 *P<0.05 for WT vs. NCX1.3 Tg. (D) EFS in the presence of atropine in longitudinal smooth
8 muscles isolated from the ileum in WT (n=3) and NCX1.3 Tg (n=4). Quantitative data on the
9
10 EFS-induced tonic contraction. Tonic contractions were expressed as percentages of 60 mM
11
12 KCl-induced contraction. *P<0.05 for WT vs. NCX1.3 Tg.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

30 **Fig. 10.** ACh- and SP-induced contraction in NCX1.3 Tg. (A) ACh-induced contractions in
31 longitudinal smooth muscles isolated from the ileum in WT (n=5) and NCX1.3 Tg (n=6).
32 Quantitative data on ACh-induced contractions. ACh-induced contractions are expressed as
33 percentages of 60 mM KCl-induced contractions. *P<0.05 for WT vs. NCX1.3 Tg. (B) t1/2
34 describes time from full amplitudes to half amplitudes of ACh-induced contraction. *P<0.05
35 for WT vs. NCX1.3 Tg. (C) SP-induced contractions in longitudinal smooth muscle strips
36 isolated from the ileum in WT (n=4) and NCX1.3 Tg (n=5). Quantitative data on SP-induced
37 contractions. SP-induced contractions are expressed as percentages of 60 mM KCl-induced
38 contractions (D) t1/2 describes the time from full amplitudes to half amplitudes of SP-induced
39 contraction. *P<0.05 for WT vs. NCX1.3 Tg.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65



Figure 1

A

NCX1

NCX2

WT



B

WT

NCX1 HET

NCX2 HET

PGP9.5







Figure 2

A



B




Figure 3
A




Figure 3

B




Figure 3

C




Figure 4





Figure 5

A



B




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10

